



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

with I deep blue turning dark violet-brown, with K the epithecium stains orange-red. This species is placed in the sect *Miscoblastia* by Dr. Zahlbruckner. On calcareous tufa near Clifton By the Sea, Los Angeles Co. Type with Dr. Zahlbruckner, cotypes in Sullivan Moss Society and Hasse herbaria.

**Rinodina euryspora** A. Zahlbr. n. sp. in litt. Thallus determinate, crustaceous, rimose-areolate, sordid yellowish brown, medium thick, areolae concave, 0.3 to 0.75 mm. wide; apothecia appressed sessile, 0.2 to 0.5 mm. wide; disk black and white pruinose or naked, thalline margin permanent, slightly elevated, entire or crenulate; epithecium continuous, sordid yellow; thecium colorless, 120 $\mu$  high, paraphyses stout, about 2 $\mu$  thick, eseptate, not furcate, tips of some slightly clavate and colored; hypothecium amber-yellow; asci inflated clavate, 8-spored; spores bluntovoid-ellipsoid, brown, bilocular, constricted, 12 $\mu$  to 15 $\mu$  long, 6 $\mu$  to 11 $\mu$  thick. Hym. gelat. stains blue with I, K—. Thallus K—, C—. On argyllaceous shale, Highland Park near Pasadena. Type with Dr. A. Zahlbruckner, cotype in herb. Hasse.

SANTA MONICA, CAL.

#### HEPATIC NOTE

The curator acknowledges with gratitude the gifts of the following specimens in response to the appeal for N. A. species in the January *BRVOLOGIST*:  
Dr. A. W. Farlow.—

*Plagiochasma Wrightii* and *Marchantia disjuncta* from Mexico. *Scapania gracilis* from Maine.

Dr. A. W. Evans.—

*Sphenobolus saxicola* from Gaspé Co., Quebec.

Miss A. Lorenz.—

*Gymnomitrium coralloides* from New Hampshire.

Collection of A. H. Brinkman from B. C.—

*Sauteria alpina*

*Scapania uliginosa*.

*Asterella Lindenberghiana*.

Mr. Severin Rapp from Sanford, Fla.—

*Metzgeria myriopoda*.

*Lejeunea spinulosa*.

*Frullania cobrensis*.

*Rectolejeunea Brittoniae*.

*Frullania Donnellii*.

*Brachiolejeunea corticalis*.

*Frullania Kunzei*.

*Leucolejeunea cuneifolia*.

*Thallocarpus Curtisii*.

*L. hanthoscarpa*.

*Sphaerocarpos texanus*.

*L. uncioloba*.

*Cephalozia floridæ*.

*Lopholejeunea Muelleriana*.

*Cephalozia Rappiana*.

*Cheilelejeunea polyantha*.

*Cololejeunea minutissima*.

*Leptocolea Jooriana*.

*C. setiloba*.

*Taxilejeunea aletusangula*.

*Displasiolejeunea Rudolphiana*.

It was an error to include the common *Scapania irrigua*. The list is now reduced to 86 species.

SUPERIOR, WIS., FEB. 1915.

GEO. H. CONKLIN.